

ON YOUNG NEUTRON STARS AS PROPELLERS AND ACCRETORS WITH CONVENTIONAL MAGNETIC FIELDS

M. Ali Alpar

Faculty of Engineering and Natural Sciences, Sabancı University, Orhanlı-Tuzla, Istanbul
81474, Turkey

Received _____; accepted _____

ABSTRACT

The similarity of rotation periods of, the anomalous X-ray pulsars (AXPs), the soft gamma ray repeaters (SGRs) and the dim isolated thermal neutron stars (DTNs) suggests a common mechanism with an asymptotic spindown phase, extending through the propeller and early accretion stages. The DTNs are interpreted as sources in the propeller stage. Their luminosities arise from frictional heating in the neutron star. If the 8.4 s rotation period of the DTN RXJ 0720.4-3125 is close to its rotational equilibrium period, the estimated propeller torque indicates a magnetic field in the 10^{12} Gauss range. The mass inflow rate onto the propeller is of the order of the accretion rates of the AXPs. The limited range of rotation periods, taken to be close to equilibrium periods, and conventional magnetic fields in the range 5×10^{11} - 5×10^{12} Gauss correspond to a range of mass inflow rates $3.2 \times 10^{14} \text{ gm s}^{-1} < \dot{M} < 4.2 \times 10^{17} \text{ gm s}^{-1}$. Observed spindown rates of the AXPs and SGRs also fit in with estimates for these magnetic fields and equilibrium periods. The source of the mass inflow is a remnant accretion disk formed as part of the fallback during the supernova explosion. These classes of sources thus represent the alternative pathways for those neutron stars that do not become radio pulsars. For the highest mass inflow rates the propeller action may support enough circumstellar material so that the optical thickness to electron scattering destroys the X-ray beaming, and the rotation period is not observable. These are the radio quiet neutron stars (RQNSs) at the centers of supernova remnants Cas A, Puppis A, RCW 103 and 296.5+10. The statistics and ages of DTNs suggest that sources in the propeller phase are quite common, maybe accounting for the majority of neutron stars formed in supernovae. AXPs are the rare cases whose $\dot{M}$ history allows them to reach the post-propeller accretion phase. This model obviates the need to postulate magnetars for AXPs

and DTNs. Frequently sampled timing observations of AXPs, SGRs and DTNs can distinguish between this explanation and the magnetar model.

Subject headings: stars:neutron–magnetic fields–accretion–supernovae:general

1. Introduction

Anomalous x-ray pulsars (AXPs) are characterized by periods in the range 6-12 secs (see Mereghetti 1999 for a review and an extensive list of references therein for properties of specific AXPs). RXJ0720.4-3125, one of the dim nearby ROSAT point sources thought to be thermally emitting neutron stars (Walter, Wolk & Neuhauser 1996, Walter & Matthews 1997) has a measured period (Haberl et al. 1997) of 8.4 s, while a recently discovered member of this class, RXJ0420.0-5022, is reported to have a period of 22.7 s (Haberl, Pietsch & Motch 1999). Among the four confirmed soft gamma ray repeaters (SGRs- Mazets et al. 1979, 1981, Cline et al. 1980, Woods et al. 1999) with quiescent X-ray sources, there are two with measured pulse periods. Both periods are in the AXP period range, with $P = 7.47$ s for SGR 1806-20 (Kouveliotou et al. 1998) and $P = 5.16$ s for SGR 1900+14 (Hurley et al. 1999, Kouveliotou et al. 1999). For the AXPs it was pointed out (van Paradijs, Taam & van der Heuvel 1995, Ghosh, Angelini & White 1995) that these periods would obtain as equilibrium periods for the accretion rates $\dot{M} \sim 10^{15} gm/s$ corresponding to the observed luminosities and for 10^{11} - 10^{12} G magnetic fields.

Values of $\dot{P}$ measured from the AXPs and SGRs, together with the long rotation periods in the 5-12 s range have led to the suggestion that these sources are magnetars (Kouveliotou et al. 1998, 1999, Duncan & Thompson 1992, Thompson & Duncan 1993, 1995), neutron stars with very strong magnetic fields $B \sim 10^{14}$ - 10^{15} G spinning down through magnetic dipole radiation. The short cooling age estimated for RXJ0720.4-3125

has also been suggested (Heyl & Hernquist 1998) as evidence for a magnetar if the source is an isolated pulsar born with a short rotation period. We propose here an alternative explanation, extending the accretion hypothesis to include propeller phases, and linking these classes of neutron stars as stars with conventional $\sim 10^{12}$ G magnetic fields. Mass inflow at rates $\dot{M} \sim 5 \times 10^{15} - 4 \times 10^{16}$ gm/s from a disk (van Paradijs, Taam & van den Heuvel 1995, Ghosh, Angelini & White 1997) around the neutron star will produce torques on the neutron star of the order of the spindown torques observed. The AXPs and SGRs are accreting at least part of the incoming mass flow. The similarity of the periods simply reflects similar conditions. All these systems are asymptotically approaching equilibrium periods in a common range, defined by the common ranges of magnetic fields and mass transfer rates. As the approach to rotational equilibrium is asymptotic, $P/\dot{P}$ is not the true age of these systems. Sources of different ages and different circumstances with similar conventional magnetic fields have similar periods in this asymptotic regime under a fairly large range of mass inflow rates.

We propose that the DTNs, AXPs, SGRs and RQNSs constitute the alternative subclasses of young neutron stars produced by supernovae, complementary to the subclass of neutron stars that show up as radio pulsars (Kaspi 1999). In view of the plausible associations of several AXPs and SGRs with supernova remnants, tight upper limits on the mass of a binary companion (Mereghetti, Israel & Stella 1998, Wilson et al. 1998), and statistics and age considerations, these sources are likely to be young neutron stars under mass inflow from a remnant disk formed by fallback material from the supernova explosion (Chevalier, 1989, Lin, Woosley & Bodenheimer 1991, Mineshige, Nomoto & Shigeyama 1993).

Most ideas discussed in this paper and their links to related subjects were first presented in an earlier preprint (Alpar 1999). The DTNs are the propeller counterparts to

the accreting AXPs. Their luminosity is due to energy dissipation in the neutron star. They may well account for a large fraction of neutron stars formed in supernovae. For the AXPs the scenario of accretion from a remnant disk, preceded with a propeller phase, has been explored also by Chatterjee, Hernquist & Narayan (1999), who have employed specific thin disk evolution models for the time dependence of the mass inflow from the remnant disk, while Marsden et al. (1999) have evaluated the supernova associations of AXPs and SGRs to argue that what unites these sources is their atypical SNR environments rather than an intrinsic property of the neutron stars like superstrong magnetic fields. Under the working hypothesis of the present work, that all supernovae form neutron stars with conventional $\sim 10^{12}$ G magnetic fields, the differences between the DTNs, AXPs and SGRs are due to their different mass inflow environments and histories. At the highest mass inflow rates we have the RQNSs. These are propellers whose pulse periods are not observable because of accumulated circumstellar material that is optically thick to electron scattering. The subclasses unified in this scenario represent the entire range of mass inflow rates, extending from $\dot{M} \sim 0$ for the radio pulsars to near Eddington rates inferred for the RQNSs.

Table 1 displays observed parameters and model estimates for the AXPs, for the two SGRs with $\dot{P}$ measurements, the two DTNs with measured periods, and for a typical RQNS. Section 2 discusses the energy dissipation in a neutron star under an external torque. Section 3 introduces propeller torques and applies these ideas to DTNs. Section 4 presents a simple model for asymptotic spindown. Section 5 relates the different classes of neutron stars, including the RQNSs, as supernova products under mass inflow. Section 6 reviews the expected signature of a remnant disk around the neutron star, and Section 7 presents the conclusions.

2. Luminosity of a Non-accreting Neutron Star from Energy Dissipation

The DTNs yield fits to blackbody spectra with temperatures of 57 eV for RXJ185635-3754 and 79 eV for RXJ0720.4-3125 and luminosities in the $L_x \sim 10^{31-32}$ erg s⁻¹ range (Walter, Wolk & Neuhauser 1996, Walter & Matthews 1997, Haberl et al. 1997, Motch & Haberl 1998, Kulkarni & van Kerkwijk 1998). Several further candidates for this class (Stocke et al. 1995, Haberl, Motch & Pietsch 1998, Schwope et al. 1999, Motch et al. 1999, Haberl, Pietsch & Motch 1999) also have similar blackbody temperatures, flux values and limits on the ratio of X-ray flux to optical flux. Accretion from the interstellar medium would require unlikely ambient interstellar medium densities and low velocities (for a discussion of accretion from the ISM for the DTNs see Treves et al. 1999 and references therein). The presence of several dim thermal neutron stars within ~ 100 pc suggests that there are $\sim 10^4$ such sources in the galaxy, requiring ages of $\sim 10^6$ years or longer, if the birth rate is 10^{-2} yr⁻¹ or less. The cooling of a young neutron star can typically yield the observed thermal luminosities at ages of the order of 10^5 - 10^6 yrs (Ögelman 1995). For a neutron star born with a rotation period of the order of 10 ms, (as typically inferred from the P and $\dot{P}$ values of young pulsars), to have spun down to the 8.4 s period of RXJ0720.4-3125 in 10^5 yrs as a rotating magnetic dipole would require a mean spindown rate $\dot{\Omega}$ of the order of 10^{-13} rad s⁻², and a magnetic field of the order of 10^{14} G or more. (Heyl & Hernquist 1998).

There is an alternative source of the thermal luminosity which takes over at $\sim 10^5$ - 10^6 yrs, after the initial cooling, and lasts longer than the cooling luminosity: There will be energy dissipation (frictional heating) in a neutron star being spun down by some external torque. The rate of energy dissipation is given by (Alpar et al. 1984, Alpar, Nandkumar & Pines 1985)

$$\dot{E}_{diss} = I_p \omega |\dot{\Omega}| \quad (1)$$

where I_p is the moment of inertia of some component of the neutron star whose rotation rate is faster than that of the observed crust by the amount ω . $\dot{E}_{diss}$ will supply the thermal luminosity of a non-accreting neutron star at ages greater than $\sim 10^6$ years (Umeda et al. 1993), as the cooling luminosity rapidly falls below $\dot{E}_{diss}$ after the transition from neutrino cooling to surface photon cooling. Among the radio pulsars with X-ray emission (Ögelman 1995, Becker & Trümper 1997), observations (Alpar et al. 1987, Yancopoulos, Hamilton & Helfand 1994) of the pulsar PSR 1929+10, whose spin-down age is 3.1×10^6 yrs, provide an upper limit to the thermal luminosity which yields $I_p\omega < 10^{43}$ gm cm² rad s⁻¹ (Alpar 1998).

A lower limit to $\dot{E}_{diss}$ can be obtained from the parameters of large pulsar glitches (Alpar 1998). The consistency of the observed glitch parameters of all pulsars with large glitches ($\Delta\Omega/\Omega \geq 10^{-7}$) and measured second derivatives $\ddot{\Omega}$ of the rotation rate (Shemar & Lyne 1996) as well as the statistics of the large glitches (Alpar & Baykal 1994) support the hypothesis that large glitches are a universal feature of pulsar dynamics. The current phenomenological description (Alpar et al. 1993, Alpar 1998) of large glitches entails angular momentum exchange between the crust and an interior component (a pinned superfluid in current models) which rotates faster than the crust by the lag ω . The typical glitch related change in relative rotation rate of the crust and interior, $\delta\omega \sim 10^{-2}$ rad s⁻², inferred from the common behaviour of all pulsars with large glitches, must be less than the lag ω . Using values of $I_p \sim 10^{43}$ gm cm² inferred from the detailed postglitch timing measurements available for the Vela pulsar glitches (Alpar et al. 1993) we obtain the lower bound $I_p\omega > I_p\delta\omega = 10^{41}$ gm cm² rad s⁻¹. This lower bound is independent of the details of the glitch models and rests only on the assumption that the large glitches involve angular momentum exchange within the neutron star. Further, as the mode of angular momentum transfer inside the star depends only on neutron star structure, the same parameter $I_p\omega$ would determine the energy dissipation rates in all neutron stars under external torques, also when the source of the external torque is not magnetic dipole radiation. The upper

and lower bounds together imply

$$10^{41}|\dot{\Omega}|ergs^{-1} < L = \dot{E}_{diss} < 10^{43}|\dot{\Omega}|ergs^{-1}. \quad (2)$$

The luminosities of RXJ0720.4-3125, RXJ185635-3754 and the other DTNs (Treves et al. 2000) give

$$|\dot{\Omega}| \sim 10^{-13} - 10^{-10} rad s^{-2}. \quad (3)$$

The measured spindown rates of AXPs and SGRs all fall in this range, providing another similarity, in addition to the periods, between the DTNs, AXPs and SGRs. AXP spindown rates actually suggest that the DTN spindown rate may be closer to $|\dot{\Omega}| \sim 10^{-12} \text{ rad s}^{-2}$, which means that $\dot{E}_{diss}$ is closer to the upper limit in Eq.(2). With the 8.4s period of RXJ0720.4-3125 these spindown rates imply surface magnetic fields in excess of 10^{14} G if magnetic dipole spindown is assumed. Are there other spindown mechanisms that will give high spindown rates, larger than $10^{-12} \text{ rad s}^{-2}$, with 10^{12} G magnetic fields typical of the canonical radio pulsars and of the accreting neutron stars with observed cyclotron lines?

3. Propeller Spindown

High spindown rates, larger than $10^{-12} \text{ rad s}^{-2}$, can indeed be expected for neutron stars with conventional 10^{12} G fields under the typical spindown torques for certain phases of accreting sources. For the AXPs accretion is a possibility that has already been explored (Mereghetti & Stella 1995, van Paradijs, Taam & van den Heuvel 1995, Ghosh, Angelini & White 1995) and will be pursued below. In connection with the DTNs, we note that a neutron star subject to mass inflow will experience high spindown rates even when the inflowing mass is not accreted, because of the star's centrifugal barrier (the propeller effect) (Illarionov & Sunyaev 1975). We propose that RXJ 0720.4-3125 and the other DTNs are

neutron stars with magnetic fields of the order of 10^{12} G, spinning down under propeller torques. There is no accretion yet in the propeller phase. The luminosities of the DTNs are produced by energy dissipation in the neutron star. These luminosities are dimmer, by several orders of magnitude, than the accretion luminosities that would have been produced if the mass inflow were accreted. The propeller torques depend on the magnetic moment of the neutron star and on the rate of mass inflow. Using the spindown rates estimated from the energy dissipation luminosities together with order of magnitude expressions for propeller torques, we show that the neutron star has a dipole magnetic field of the order of 10^{12} G. The estimated range of mass inflow rates coincides with the range of mass accretion rates inferred from the X-ray luminosities of the accreting AXPs and SGRs, suggesting that the DTNs are in the propeller stage and the AXPs and SGRs in the accretion phase under similar mass inflow circumstances.

Occasional quiescent states of some X-ray transients are likely examples of the propeller phase (Cui et al 1997, Zhang et al 1998, Campana et al 1998, Menou et al. 1999). Sources that are persistently in the propeller phase have not been detected previously. This is understandable since they are not lit up with an accretion luminosity. Thus RXJ 0720.4-3125, RXJ185635-3754 and other DTNs are the first observed examples of propellers, observed only through their dissipation luminosities, only as the nearest members of the DTN class, and only by ROSAT, in view of their surface temperatures in the soft X-ray band.

In the propeller phase there is a spindown torque exerted on the neutron star, through the interaction of its magnetosphere with the ambient material. The order of magnitude of this spindown torque is

$$N = I|\dot{\Omega}| = \mu^2/r_A^3 \quad (4)$$

where $\mu = BR^3$ is the magnetic moment of the neutron star with surface dipole magnetic

field B and radius R , and

$$r_A = 9.85 \times 10^8 cm \mu_{30}^{4/7} \dot{M}_{15}^{-2/7} m^{-1/7} \quad (5)$$

is the Alfven radius (Lipunov 1992, Frank, King & Raine 1992). Here μ_{30} denotes the magnetic moment in units of 10^{30} G cm³, $\dot{M}_{15}$ is the mass inflow rate in units of 10^{15} gm s⁻¹ and m is the neutron star mass in solar mass units. The propeller phase will last until accretion starts when a critical rotation period is reached. This critical period is of the order of, but somewhat smaller than the equilibrium rotation period

$$P_{eq} = 16.8 s \mu_{30}^{6/7} \dot{M}_{15}^{-3/7} m^{-5/7} \quad (6)$$

which obtains when the star's rotation rate equals the Keplerean rotation rate of ambient matter at the Alfven radius; that is, when the corotation radius $r_c = (GM)^{1/3} \Omega^{-2/3}$ becomes equal to r_A . In the propeller phase the source settles to an asymptotic spindown towards the equilibrium. A source is most likely to be observed during its asymptotic phase. Since in the asymptotic regime r_c is close to r_A we can estimate the magnetic moment of a propeller from its spindown rate, and rotation rate, without knowing the mass inflow rate, simply by substituting r_c in Eq.(4). This leads to

$$\mu = \frac{(I|\dot{\Omega}|GM)^{1/2}}{\Omega}. \quad (7)$$

We take the DTN RXJ0720.4-3125, with $P = 8.4$ s, as our typical example (Haberl et al. 1997). The other DTN with a reported rotation period, RXJ 0420.0-5022, $P = 22.7$ s (Haberl, Pietsch & Motch 1999), yields similar results. For RXJ0720.4-3125 we obtain $\mu_{30} \sim (0.35 - 3.5)$ using the bounds on the spindown rate, Eq.(2). Thus if RXJ0720.4-3125 is a propeller it has a conventional magnetic field,

$$3.5 \times 10^{11} G < B < 3.5 \times 10^{12} G \quad (8)$$

Using Eq.(6) and setting the observed period to be equal to the equilibrium period, we can now obtain bounds on the mass inflow rates:

$$\dot{M}_{15} = (16.8s/P_{eq})^{7/3}(\mu_{30})^2 m^{-5/3}. \quad (9)$$

This yields $\dot{M}_{15} \sim (0.62 - 62)$ for the mass inflow rate for RXJ0720.4-3125. Model parameters obtained from Eqs. (7) and (9) are given in Table 1, assuming $|\dot{\Omega}| = 2.6 \times 10^{-12}$ rad s⁻² from Eq.(2) for RXJ0720.4-3125, and $|\dot{\Omega}| = 10^{-11}$ rad s⁻² for RXJ 0420.0-5022.

For most DTNs and for all RQNSs, which, as we propose below, are also propeller candidates, we do not have observed rotation periods. For these sources, setting the rotation period to be equal to the equilibrium period,

$$\dot{M}_{15} \cong \mu_{30}^{-1/3} |\dot{\Omega}|_{-12}^{7/6} m^{-1/2} I_{45}^{7/6} \quad (10)$$

from Eqs.(6) and (7). With the range of $|\dot{\Omega}|$ obtained from DTN luminosities (Eqn. (3)), $\dot{M}_{15} \sim (0.1 - 200)$ is obtained for these DTNs for $\mu_{30} = 1$.

If the source of mass inflow is not depleted, the propeller phase is followed by an accretion phase starting at some critical period. The neutron star will continue to spin down, now as an accreting source, as its period evolves from the critical period towards the equilibrium period. Similarity of the rotation periods of the AXPs with the 8.4 s period of the DTN RXJ0720.4-3125 suggests that the approach to rotational equilibrium is an asymptotic process extending through the propeller phase on to the accretion phase. These sources are asymptotically close to their individual equilibrium rotation periods, which lie in a narrow range determined by their magnetic moments and mass inflow histories.

The mass accretion rates inferred from the X-ray luminosities of the AXPs and SGRs all fall in the range $\dot{M}_{15} \sim 1 - 100$. The corresponding magnetic moments for accreting sources near rotational equilibrium are in the 10¹² G range given the 5-15 s rotation periods:

$$\mu_{30} = (P_{eq}/16.8s)^{7/6} \dot{M}_{15}^{1/2} m^{5/6}. \quad (11)$$

To summarize, (i) the spindown rates inferred for the DTN sources through the interpretation of their luminosities as due to energy dissipation (Eqs.(1)-(3)) agree with the observed spindown rates for the AXPs and SGRs to order of magnitude. (ii) Estimates of the magnetic moment, from the spindown rates for the DTNs as propellers close to rotational equilibrium, coincide in the same range as the magnetic moments inferred from the mass accretion rates implied by the X-ray luminosities of AXPs and SGRs as accretors asymptotically close to rotational equilibrium. This is the conventional 10^{12} G range of most young neutron stars observed as rotation powered pulsars and in high mass X-ray binaries. (iii) Furthermore, the mass inflow rates inferred for the DTNs from the assumption that they are close to rotational equilibrium, agree with the accretion rates of the AXPs and SGRs. These observations strongly support the unifying hypothesis that the sources are related instances of asymptotic spindown near similar equilibrium periods.

4. The Asymptotic Spindown Regime

We now turn to a simple model for the asymptotic spindown. A neutron star in the presence of inflowing matter experiences both spindown and spin-up torques. The long term evolution is determined by the balance between these, described by a function $n(\Omega/\Omega_{eq} - \omega_c)$ (Ghosh & Lamb 1991) which goes through zero when $\Omega = \omega_c \Omega_{eq}$ where ω_c is of order one, and $\Omega_{eq} = 2\pi/P_{eq}$. In accretion from a disk, the relative specific angular momentum brought in by the accreting material to spin the neutron star up is $[(GM r_A)^{1/2} - \Omega r_A^2]$. Since the dimensional torque is $\mu^2/r_A^3 \sim \dot{M}(GM r_A)^{1/2}$, spindown near equilibrium can be modelled as

$$\dot{\Omega} = (\mu^2/I r_A^3)(1 - \Omega/\Omega_{eq}) = (\Omega_{eq} - \Omega)/t_0 \quad (12)$$

where $t_0 = \Omega_{eq} I r_A^3 / \mu^2$. Here the zero of the torque (the end of the spindown era) is taken to be at $\Omega = \Omega_{eq}$ rather than $\omega_c \Omega_{eq}$, for simplicity. The AXPs and SGRs must then

be sources that have emerged from propeller spindown and started to accrete while still under a net spindown torque as they have not yet reached rotational equilibrium. Another simplifying assumption we make is that the mass inflow rate is constant in each source. In a more realistic model the mass inflow rate should be time dependent, especially if its source is a remnant disk with a finite lifetime, rather than a binary companion. A decaying $\dot{M}$ leads to a natural explanation for the prevalence of spindown in an accreting source: the equilibrium period increases as $\dot{M}$ decreases, and the source will be spinning down as its period tracks the equilibrium period (Ghosh, Angelini & White 1997, Chatterjee, Hernquist & Narayan 1999), answering the criticism of accretion models raised by Xi (1999). Chatterjee, Hernquist & Narayan (1999) have employed an $\dot{M}$ decaying as a power law in time, corresponding to the viscous evolution of a thin disk (Mineshige, Nomoto & Shigeyama 1993, Canizzo et al 1997). The actual situation warrants taking into account propeller boundary conditions. The propeller activity could probably support a thick disk or corona. It would also lead to a different characteristic time dependence of $\dot{M}$. Here we take a constant $\dot{M}$ to represent an average of the actual $\dot{M}(t)$.

Even under constant mass inflow rate, the torque history must be more complicated. In X-ray binaries, the occurrence of spindown and spinup torques of comparable magnitudes, and torque reversals on all timescales signals the importance of electromagnetic torques. The real torques could lead to initial power law decays of Ω as a function of time. Whatever the form of the initial spindown may be, once the rotation rate is close to the equilibrium value, the factor $(\Omega_{eq} - \Omega)$ dominates the asymptotic evolution, which becomes an exponential decay. Here we employ the simple asymptotic model of Eq.(12) assuming that the magnetic moment is constant, and that the mass inflow rate is also a constant representing the long term average $\dot{M}$. The results are given in Table 1 and Figure 1. With constant μ and r_A , the spindown leads Ω towards Ω_{eq} :

$$\Omega(t) = [\Omega(0) - \Omega_{eq}]exp(-t/t_0) + \Omega_{eq}. \quad (13)$$

This solution is to describe the spindown through both the propeller and the accretion phases, with mass accretion starting at some $\Omega_{acc} < \Omega_{eq}$. From Eqs.(5),(6) and (12) we obtain

$$t_0 = 1.3 \times 10^{12} s I_{45} \dot{M}_{15}^{-1} m^{-2/3} \Omega_{eq}^{4/3}. \quad (14)$$

Substituting the expression for t_0 in the spindown equation, Eq.(12), Ω_{eq} can be obtained for each source with known Ω and $\dot{\Omega}$ by solving

$$\Omega_{eq} = \Omega + 1.3 \times 10^{12} s \dot{\Omega} I_{45} \dot{M}_{15}^{-1} m^{-2/3} \Omega_{eq}^{4/3}. \quad (15)$$

Once the equilibrium rotation rate is estimated, the magnetic moment μ can be obtained from Eq.(11). The time since the beginning of the propeller phase can be obtained from Eq.(13) for a given initial rotation period, which can be taken to be $P(0) = 0.01$ s for a newborn neutron star. The age t estimated from Eq.(13) is:

$$t = t_0 \log[(\Omega(0) - \Omega_{eq})/(\Omega - \Omega_{eq})]. \quad (16)$$

Sample solutions are given in Table 1 for the AXPs, the SGRs with measured P and $\dot{P}$ and for the DTNs with measured periods. The AXPs show changes in spindown rate, by up to an order of magnitude, on timescales of several years. Long term average values of the spindown rate are used, as appropriate for the model to describe the long term average trend. Mass inflow rates are inferred from the observed luminosities taken as accretion luminosities onto neutron stars, $\dot{M}_{acc} = RL/GM$. The luminosity has been observed to change by as much as a factor of 15 in AXPs (Torii et al. 1998). Furthermore, the mass inflow rate from a remnant disk will be decreasing as a function of time, so that the constant representative $\dot{M}$ appropriate for our simple model should be larger than the present time $\dot{M}_{acc}$ inferred from the accretion luminosity. For 1E1048.1-5937, the current $\dot{M}_{acc}$ leads to $P_{eq} = 64$ s., while for 1E2259+586 the current $\dot{M}_{acc}$ gives a relaxation time t_0 that may be larger than the age of the associated SNR. For these two sources, taking examples of mass

inflow rates $\dot{M} > \dot{M}_{acc}$, one can obtain solutions in agreement with the asymptotic model. These solutions are given in Table 1.

Three out of six AXPs provide direct evidence of youth through their likely supernova associations. These AXPs are situated close to the centers of the SNRs. The associations with SNRs are not as certain for the SGRs: The SGRs are at edges of the SNRs, requiring velocities $\sim 1000 \text{ Km s}^{-1}$ if they were born at the center (for a recent discussion of the SNR associations of the AXPs and SGRs, and references to the individual associations, see Gaensler 1999). For both AXPs and SGRs, the associated SNRs are not typical; but rather of a particularly dense morphology. This point was realized and elaborated by Marsden et al (1999) in their discussion of propeller and accretion scenarios, as evidence that these classes of sources are related through similarities in their environments rather than a common intrinsic property like very strong magnetic fields. Here we adopt the possibility that a remnant disk around the neutron star, formed from the debris of the core collapse in a supernova explosion (Chevalier 1989, Mineshige, Nomoto & Shigeyama 1993), provides the mass inflow. The propeller and accretion phases would terminate when the supply of $\dot{M}$ is depleted.

5. The Fallback Mass Inflow and the Signature of the Neutron Star Born in a Supernova

Does the narrow range of observed rotation periods indicate a very restricted range of $\dot{M}$? The range of estimates for the sources in Table 1 extends from $10^{15} \text{ gm s}^{-1}$ to $10^{17} \text{ gm s}^{-1}$. To estimate the parameter range corresponding to the narrow range of equilibrium periods, we note that the magnetic moment and mass inflow rate or equivalently, the magnetic moment and equilibrium rotation rate are independent parameters. For magnetic moments in the range $5 \times 10^{11} \text{ G} - 5 \times 10^{12} \text{ G}$, the range of mass inflow or accretion rates

which would lead to equilibrium periods of 5-15 s extends over three orders of magnitude, up to about half the Eddington rate:

$$3.2 \times 10^{14} \text{ gms}^{-1} < \dot{M} < 4.2 \times 10^{17} \text{ gms}^{-1}. \quad (17)$$

This is not a very restricted range. Fig.1 shows the mass inflow rate as a function of the equilibrium period for $0.5 < \mu_{30} < 5$. Also shown are the model solutions for equilibrium periods and mass inflow rates for all the sources presented in Table 1. The narrow range of observed periods lies between the vertical lines. We note that the narrow range of observed and equilibrium periods corresponds to a rather wide range of $\dot{M}$, extending over more than two orders of magnitude. Sample solutions for different sources are scattered in $\dot{M}$ and μ , as they should be, since the magnetic moments of neutron stars should not be correlated with the mass inflow from their environments. This scatter indicates that the model does not require an artificial correlation between these parameters.

The mass inflow rates we obtain are all above $\sim 10^{14} \text{ gm s}^{-1}$. This means that for lower $\dot{M}$ the propeller phase never starts, and the neutron star promptly starts life as a radio pulsar. At the other extreme extended exposure to the highest mass inflow rates may lead to accumulation of enough circumstellar material such that the optical thickness to electron scattering washes away any beaming of the X-rays from the neutron star and the rotation period is not detected (Lamb et al 1985, Ghosh, Angelini & White 1997).

If all supernovae left neutron stars that went through an AXP phase under mass inflow from the debris, then we would expect at least 100 such objects for a lifetime $> 10^4$ yrs and galactic supernova rate of 10^{-2} , since the AXPs would be observable from all galactic distances. That we only observe a few indicates that such sources rarely reach the accretion phase.

The number of DTNs must be much larger, since we see several within a distance of the order of 100 pc. In the Galaxy we would expect about 10^4 DTNs, and a lifetime of

the order of 10^6 yrs or longer, for a birth rate of 10^{-2} yr^{-1} or lower. The much smaller number of the AXPs may be due to the depletion of the supply of mass inflow in most sources by the end of the propeller phase. If the mass inflow is depleted before the end of the propeller phase, subsequent accretion phases never occur. This may be the reason why the numbers of AXPs and SGRs are much less than the number of DTNs. By the same token, mass inflow rates $< 3.2 \times 10^{14} \text{ gm s}^{-1}$ are not inferred: at lower mass inflow rates the duration of the propeller phase exceeds the lifetime of the mass inflow source. The rare AXPs and SGRs are thus the young objects born with conventional magnetic fields but in circumstances of large enough mass inflow towards the neutron star, so that the time t_0 , which scales with $\dot{M}^{-1}$, is short compared to the lifetime of the mass supply: These are the rare sources which have gone through the propeller phase in time to start accretion before the matter supply is depleted. The more common source is the DTN, in a propeller phase of duration 10^6 yrs, surviving the disappearance of the supernova remnant. While it is reasonable to expect that the lifetime of the mass supply increases at the lower mass inflow rates, one needs a model of the disk evolution to make the comparison with the propeller spindown quantitative. If DTNs are born at a rate comparable to the total supernova rate, then a significant fraction of SNRs must leave the neutron star under conditions of low enough $\dot{M}$. If all AXPs are to be descendants of DTNs, either the propeller phase lasts 1000 times longer than the AXP accretion phase, or only 10^{-3} of DTNs turn into AXPs, according to the comparison of total numbers of these two types of sources in the Galaxy. Taking into account the evidence for possible association with supernova remnants, and the agreement between SNR ages and the age estimates the asymptotic model, mass supply from a debris disk around a single neutron star seems to be the likely scenario, rather than mass supply from a binary companion.

The SGRs are like the AXPs in their X-ray properties. Both classes of sources are detectable throughout the galaxy, and their comparable numbers suggest the SGRs are also

in the same rare or relatively short beginning accretor phase as the AXPs. The claimed associations of most SGRs with SNRs are less certain than the AXP-SNR associations. Unlike the AXPs the SGRs are located at the outskirts of their candidate SNR associations. Nevertheless these SNRs are of the same atypical dense class as the SNRs associated with the AXPs (Marsden et al 1999).

For the SGRs the magnetar hypothesis is on a different, stronger footing than is the case for the AXPs, in that it provides the energy store and a detailed dynamical model for the soft gamma ray bursts. While the present work does not address the mechanisms of the gamma ray burst phenomenon, it is nevertheless intriguing that as Wang and Robertson (1985) have noted propellers (and probably their descendants the early accretors) can support relativistic particle luminosities and gamma ray production in the surrounding accumulated matter. The energy source could be sporadic accretion of accumulated circumstellar mass released through instabilities. The remnant disk scenarios were first recalled in connection with the problem of planet formation around pulsars (Lin, Woosley & Bodenheimer 1991), and disk instabilities involving planet-like masses (Mineshige, Nomoto & Shigeyama 1993) are consistent with the luminosities of the SGRs. It may not be unreasonable to speculate that SGRs are also a class of neutron stars with conventional fields undergoing disk instabilities.

Why is it that the AXPs are observed as pulsars while the LMXBs are not ? The important single exception is the recently discovered source XTE J1808-369. The explanation for the rarity of LMXB pulsars has been that comptonization by circumstellar material will destroy pulses by washing out the beams emerging from the neutron star if the material is optically thick to electron scattering (Lamb et al. 1985). This was noted in connection with the AXPs by Ghosh, Angelini & White (1997). The present picture is consistent with this: the AXPs are observed as pulsars because they are in the beginning

stages of accretion, and the corona around them does not have significant optical thickness $\tau_{es} > 1$ to destroy the beaming. Work in progress to test this hypothesis by modelling the AXP spectra as unsaturated comptonization and estimating the optical thicknesses will be reported separately.

The radio quiet neutron stars found in centers of the supernova remnants Cas A, Puppis A, RCW 103 and 296.5+10 (Chakrabarty et al. 2000, Gaensler, Bock & Stappers 2000, Petre, Becker & Winkler 1996, Gotthelf, Petre & Hwang, 1997, Gotthelf, Petre & Vasisht 1999, Mereghetti, Bignami & Caraveo 1996, Vasisht et al. 1997) may be the propeller sources for which the cumulative effect of the mass inflow has indeed set up a corona of $\tau_{es} > 1$ around the neutron star. The source properties are very similar. Variability of the flux (Gotthelf, Petre & Vasisht 1999) favours accretion or propeller driven energy dissipation rather than the cooling luminosity of an isolated neutron star. The low luminosities, $L_x \sim 10^{33} - 10^{34} \text{ erg s}^{-1}$ suggest that these sources are in the propeller phase. Accretion would imply $\dot{M} \sim 10^{13} - 10^{14} \text{ gm/s}$, which can accrete onto the neutron star only if the rotation period is longer than 50 s (Eq.(6)) for $B \sim 10^{12} \text{ G}$. But then at these low accretion rates there is no reason that we should not see pulses at the rotation period. Blackbody effective areas of 1 km^2 also enhance the expectation of coherent pulsations at the rotation period from these sources while upper limits of 13% and 25% have been obtained for the RQNSs in RCW103 and Cas A respectively (Gotthelf, Petre & Vasisht 1999, Chakrabarty et al. 2000). Taking the RQNSs to be propeller sources with $L_x = \dot{E}_{diss}$, as we did for the DTNs, leads to $|\dot{\Omega}| \sim 10^{-10} - 10^{-8} \text{ rad s}^{-2}$. Since the rotation periods of the RQNSs are not known, we use Eq.(10) with $B \sim 10^{12} \text{ G}$, and obtain

$$\dot{M} > 2 \times 10^{17} \text{ gm s}^{-1}. \quad (18)$$

For the RQNSs the mass inflow rates $\dot{M}$ are indeed higher than those inferred for the DTNs and AXPs, consistent with the proposal that the radio quiet neutron stars are the high

$\dot{M}$ end of our spectrum of neutron stars under mass inflow from remnant disks. Model parameters for the RQNS in Cas A are displayed in Table 1. The luminosity is typical of the RQNSs. As a propeller with luminosity arising from energy dissipation, taking $|\dot{\Omega}| > 3 \times 10^{-10} \text{ rad s}^{-2}$ with $\mu_{30} = 1$ and obtain $\dot{M} > 7.4 \times 10^{17} \text{ gm s}^{-1}$, in agreement with the interpretation here. $P_{eq} < 1 \text{ s}$, corresponding to the high mass inflow rate. The unobserved periods of the RQNSs are smaller than the observed range of periods of the AXPs and DTNs.

6. Luminosity of the Disk

For both the DTNs and the AXPs there will be another source of energy dissipation in the disk or circumstellar material, due to accretion down to r_A :

$$L(r_A) = 3/2 GM \dot{M} / r_A \sim 3/2 (GM)^{2/3} \Omega^{2/3} \dot{M} \sim 1.3 \times 10^{33} \text{ erg s}^{-1} \dot{M}_{15} m^{2/3} P^{-2/3} \quad (19)$$

Near rotational equilibrium, $L(r_A) = I \Omega \dot{\Omega}$, the power expended on the star by the spindown torque. For the AXPs this luminosity is smaller than the accretion luminosity by a factor

$$L(r_A)/L \sim 3/2 R/r_A (\dot{M}/\dot{M}_{acc}) \sim 10^{-2} (\dot{M}/\dot{M}_{acc}) m^{-1/3} P^{-2/3} \quad (20)$$

noting that the mass inflow may not be all accreted even in the AXPs. For the DTNs $L(r_A)$ is actually larger than the observed luminosity $\dot{E}_{diss}$ by a factor

$$L(r_A)/\dot{E}_{diss} \sim (I \Omega |\dot{\Omega}|) / (I_p \omega |\dot{\Omega}|) \sim (10^2 - 10^4) \Omega. \quad (21)$$

If this luminosity is dissipated entirely at the boundary region, $r \sim r_A$, the effective temperature is:

$$T(r_A) = (L(r_A)/4\pi r_A^2 \sigma)^{1/4} \sim 9.5 \times 10^4 K \dot{M}_{15}^{1/4} P^{-1/2} \quad (22)$$

If the disk temperature lies in the euV range this would be extremely difficult to detect, but being nearby sources, DTNs with low neutral hydrogen column density might provide

a chance of looking for a disk luminosity as a test of the present model. For the thin disk model employed by Chatterjee, Hernquist & Narayan (1999) calculations of the expected disk spectrum and luminosity are reported by Perna, Hernquist & Narayan (2000). For $T(r_A)$ in the optical or IR, observational upper limits for disks around the AXPs (Coe and Pightling 1998, Hulleman et al 1999) pose a problem for thin disk models. The upper limits allow thin disks with disk outer radii $\sim 10^{10}$ cm, which requires a very young age, $t \sim 100$ yrs, in terms of the viscous evolution of an isolated thin disk. However several considerations suggest that the remnant disks may be consistent with the observational upper bounds. The propeller effect might well lead to boundary conditions that alter the disk evolution. The work done on the inflowing material by the propeller, at the rate $I\Omega\dot{\Omega}$, may lift and sustain the material at distances much larger than r_A . The emerging radiation may have a lower effective temperature, from a larger effective area. The remnant disk may be a thick disk, and it may be enshrouded in a comptonizing corona, as we invoked above. Modelling of remnant disks and circumstellar material with propeller boundary conditions, and future observations on the spectra of DTNs, AXPs and radio quiet neutron stars will help to resolve these issues.

7. Discussion and Conclusions

A unified picture is proposed to account for all neutron stars formed in supernovae, and to include RQNSs, DTNs and AXPs and perhaps SGRs. The salient features of this picture are:

(i) The signature of a young neutron star depends on the presence and nature of the mass inflow of fallback material from the supernova explosion. The related classes of sources represent different pathways under different mass inflow rates and histories.

(ii) All neutron stars are born with 10^{12} G fields.

(iii) Radio pulsars are formed if there is no mass inflow or not enough to protrude the light cylinder. Higher mass inflow rates do not allow radio pulsar activity, leading to DTNs, AXPs and RQNSs.

(iv) The similarity in the rotation periods of these sources is not a coincidence but rather a consequence of the asymptotic approach to rotational equilibrium under a wide range of mass inflow rates.

(v) With low $\dot{M}$ the spindown is too slow for the source to go through the propeller stage and reach the accretion stage before the circumstellar mass is depleted. These sources, observed as DTNs, are propellers for as long as 10^5 - 10^6 yrs, and make up the most numerous class.

(vi) The DTNs are the first observed examples of neutron stars in the propeller phase.

(vii) A non-accreting neutron star under propeller spindown has a luminosity provided by energy dissipation inside the star.

(viii) The observability of the rotation period in the AXPs but not in the LMXBs can be explained qualitatively in terms of comptonization as supported by an interpretation of their spectra. The RQNSs are the young neutron stars for which comptonization washes out beaming so that the rotation period is not observable.

(vi) With low $\dot{M}$ the spindown is too slow for the source to go through the propeller stage and reach the accretion stage before the circumstellar mass is depleted. These sources, observed as DTNs, are propellers for as long as 10^5 - 10^6 yrs, and make up the most numerous class.

(vii) The DTNs are the first observed examples of neutron stars in the propeller phase.

(viii) A non-accreting neutron star under propeller spindown has a luminosity provided by energy dissipation inside the star.

Arguments that AXPs and DTNs are magnetars are based on the grounds that isolated neutron stars with ordinary 10^{12} G fields cannot have spun down to ~ 10 s periods within the estimated ages. The above picture obviates the need to postulate magnetars for AXPs and DTNs. For the SGRs the required energy budgets and dynamical arguments make a case for the magnetar hypothesis. Marsden et al (1999) have noted that AXPs and SGRs share a common morphology of their associated supernova remnants, which they link with the high density of the surrounding interstellar medium. These authors argue that the common properties of these classes of sources must therefore be due to a common set of environmental factors, rather than intrinsic neutron star properties like superstrong magnetic fields. Whether the large spindown rates observed are due to spindown by interaction with ambient matter, as proposed here, as well as by Chatterjee, Hernquist & Narayan (1999) and Marsden et al. (1999) or to spindown by a magnetar can be decided by detailed analysis of the fluctuations (noise) in the spindown process, as the timing noise characteristic of accretion powered neutron stars is quite distinguishable from the timing noise in the typically much quieter isolated rotation powered pulsars. This analysis will require frequently sampled timing observations of the AXPs and SGRs. The recently reported extended quiet spindown phases in 1E2259+586 and 1RXS J170849.0-400910 (Kaspi, Chakrabarty & Steinberger 1999) is not sufficient to conclude that this source is undergoing spindown under magnetic dipole radiation since the quiet spindown phase was preceded by episodes of higher spindown rates with large timing noise strengths (Baykal & Swank 1996). Whether such changes in the torque repeat periodically as foreseen for a precessing magnetar (Melatos 1999) remains to be checked in future timing observations. A well known accreting source, the LMXB 4U1626-67, has exhibited similar quiet episodes as well as intervals of strong timing fluctuations that are typical of accreting

sources (Chakrabarty et al. 1997). It is encouraging from the unified point of view of this work that during the quiet spindown epoch that would allow the detection of glitches, 1E 2259+586 has indeed exhibited a glitch that is very similar to the glitches of the Vela pulsar and other radio pulsars (Kaspi, Lackey & Chakrabarty 2000). If the AXPs are confirmed as accreting sources, the glitch from 1E2259+586 will constitute strong support for our starting hypothesis that all neutron stars have the same internal dynamics and the associated energy dissipation rates. Observation of a spindown rate in the expected range from RXJ0720.4-3125 or periods and spindown rates from other DTNs would constitute strong evidence for the propeller hypothesis.

The framework proposed here leads to a program of related issues to be reported in subsequent work. These include models of the disk and circumstellar material, the emerging spectrum and pulse content, the time evolution of the mass inflow and the resulting spindown.

I thank H. Ögelman, Ş.Balman, A.Baykal, A.Esendemir, O. Guseinov, Ü.Kızıloğlu and members of the High Energy Astrophysics Research Unit at METU for useful comments and Ç. İnəm for help with the table and the figure. I acknowledge partial support from the Scientific and Technical Research Council of Turkey (TÜBİTAK), from the Turkish Academy of Sciences and from the NSF International Grant 9417296 at the University of Wisconsin-Madison.

REFERENCES

- Alpar,M.A. 1998, in Neutron Stars and Pulsars, N. Shibazaki, N. Kawai, S. Shibata & T. Kifune, eds., (Universal Academic Publishers: Tokyo), 129
- Alpar, M.A. 1999, <http://xxx.lanl.gov/abs/astro-ph/9912228>
- Alpar,M.A. & Baykal,A. 1994, MNRAS, 269, 849
- Alpar,M.A. et al. 1984, ApJ, 276, 325
- Alpar,M.A. et al. 1987, A&A, 276, 101
- Alpar,M.A., Nandkumar,R. & Pines,D. 1985, 288, 191
- Alpar,M.A., Chau,H.F., Cheng,K.S. & Pines,D. 1993, ApJ, 409, 345
- Baykal,A. & Swank,J.H. 1996, ApJ, 460, 470
- Becker,W. & Trümper,J. 1997, A&A, 326, 682
- Brazier, K.T.S. & Johnston,S. 1999, MNRAS, 305, 671
- Campana,S. et al. 1998, ApJ, 499, L65
- Canizzo,J.K., Lee,H.M. & Goodman,J. 1990, ApJ, 351,38
- Chakrabarty,D. et al. 1997, ApJ, 474, 414
- Chakrabarty,D. et al. 2000, <http://xxx.lanl.gov/abs/astro-ph/0001026>
- Chatterjee,P., Hernquist, L. & Narayan,R. 1999, <http://xxx.lanl.gov/abs/astro-ph/9912137>
- Chevalier,R.A. 1989, ApJ, 346, 847
- Cline,T. et al. 1980, ApJ, 237, L1
- Coe,M.J. & Pightling,S.L. 1998, MNRAS, 299,233
- Cui,W. et al. 1998, ApJ, 502, L49
- Duncan,R.C. & Thompson,C. 1992, ApJ 392, L9
- Frank,J., King,A. & Raine,D. 1992, Accretion Power in Astrophysics, Cambridge University Press
- Gaensler,B.M. 1999, <http://xxx.lanl.gov/abs/astro-ph/9911190>
- Gaensler,B.M., Bock,D.C.-J. & Stappers,B.W. 2000, <http://xxx.lanl.gov/abs/astro->

ph/0003032

Gotthelf, E.V., Petre, R. & Hwang, U. 1997, *ApJ*, 487, L175

Gotthelf, E.V., Petre, R. & Vasisht, G. 1999, <http://xxx.lanl.gov/abs/astro-ph/9901371>

Ghosh, P. & Lamb, F.K. 1991, in *Neutron Stars: Theory and Observation*, J. Ventura & D. Pines, eds. (Kluwer:Dordrecht), 363

Ghosh, P., Angelini, L. & White, N.E. 1997, *ApJ*, 478, 713

Haberl, F. et al. 1997, *A&A*, 326, 662

Haberl, F., Motch, C. & Pietsch, W. 1998, *Astron. Nachr.*, 319, 97

Haberl, F., Pietsch, W. & Motch, C. 1999, <http://xxx.lanl.gov/abs/astro-ph/9911159>

Heyl, J.S. & Hernquist, L. 1998, *MNRAS* 297, L69

Hulleman, F. et al. 2000, <http://xxx.lanl.gov/abs/astro-ph/0002474>

Hurley, K. et al. 1999, *ApJ* 510, L111

Illarionov, A.F. & Sunyaev, R.A. 1975, *A&A*, 39, 185

Kaspi, V.M. 1999, <http://xxx.lanl.gov/abs/astro-ph/9912284>

Kaspi, V.M., Chakrabarty, D. & Steinberger, J. 1999, <http://xxx.lanl.gov/abs/astro-ph/9909283>

Kaspi, V.M., Lackey, J.R. & Chakrabarty, D. 2000, *ApJ*, submitted

Kouveliotou, C. et al. 1998, *Nature* 393, 235

Kouveliotou, C. et al. 1999, *ApJ* 510, L115

Kulkarni, S.R. & van Kerkwijk, M.H. 1998, *ApJ*, 507, L49

Lamb, F.K., Shibazaki, N., Alpar, M.A. & Shaham, J. 1985, *Nature*, 317, 681

Lin, D.N.C., Woosley, S.E. & Bodenheimer, P.H. 1991, *Nature*, 353, 827

Lipunov, V.M. 1992, *Astrophysics of Neutron Stars*, (Springer)

Marsden et al. 1999

Marsden, D., Lingefelter, R.E., Rothschild, R.E. & Higdon, J.C. 1999, <http://xxx.lanl.gov/abs/astro-ph/9912207>

- Mazets, E.P. et al. 1979, *Nature* 282, 587
- Mazets, E.P. et al 1981, *Ap&SS* 80, 3
- Melatos,A. 1999, *ApJ*, 519, L77
- Menou et al. 1999, *ApJ*, 520, 276
- Mereghetti, S. 1995, *ApJ*, 442, L17
- Mereghetti, S. 1999, <http://xxx.lanl.gov/abs/astro-ph/9911252>
- Mereghetti, S. & Stella, L. 1995, *ApJ*, 442, L17
- Mereghetti, S., Bignami, G.F. & Caraveo, P.A. 1996, *ApJ*, 464, 842
- Mereghetti, S., Israel, G.L. & Stella, L. 1998, *MNRAS*, 296, 698
- Mineshige,S., Nomoto,K. & Shigeyama,T. 1993, *A&A*, 267, 95
- Motch, C. & Haberl, F. 1998, *A&A*, 333, L59
- Motch, C. et al. 1999, *A&A*, in the press, [astro-ph/9907306](http://xxx.lanl.gov/abs/astro-ph/9907306)
- Ögelman,H. 1995, in *The Lives of the Neutron Stars*, M.A. Alpar, Ü. Kızıloğlu & J. van Paradijs eds., (Kluwer:Dordrecht), 101
- Perna,R., Hernquist,L. & Narayan, R. 2000, *ApJ*, submitted
- Petre, R., Becker, C.M. & Winkler, P.F. 1996, *ApJ*, 465, L43
- Schwope et al. 1999, *A&A*, 341, L51
- Shemar,S.L. & Lyne,A.G. 1996, *MNRAS*, 282, 677
- Stoeck, J.T. et al. 1995, *AJ*, 109, 1199
- Thompson, C. & Duncan, R.C. 1993, *ApJ* 408, 194
- Thompson, C. & Duncan, R.C. 1995, *MNRAS* 275, 255
- Torii, K. et al. 1998, *ApJ*, 503, 843
- Treves et al. 1999, <http://xxx.lanl.gov/abs/astro-ph/9911430>
- Umeda,H. et al. 1993, *ApJ*, 408, 186
- van Paradijs,J.,Taam, R.E.& van den Heuvel,E.P.J. 1995, *A&A* 299, L41
- Vasisht, G. et al. 1997, *ApJ*, 476, L43

- Walter,F.M.,Wolk,S.J.& Neuhauser,R. 1996, Nature 379, 233
- Walter,F.M.&Matthews,L.D. 1997, Nature 389, 358
- Wang,Y.-M. & Robertson,J.A. 1985, A&A, 151, 361
- Wilson, C.A. et al. 1998, ApJ 513, 464
- Woods, P.M. et al. 1999, ApJ 519, L139
- Xi,X.-D. 1999, ApJ, 520, 271
- Yancopoulos,S., Hamilton,T.T. & Helfand,D.J. 1994, ApJ, 429, 832
- Zhang,S.N., Yu,W. & Zhang,W. 1998, ApJ, 494, L71




Fig. 1.— Equilibrium periods, in seconds, and mass inflow rates, in $10^{15} \text{ gm s}^{-1}$, of the sources in Table 1. The curves mark $\mu_{30} = 5$ (upper) and $\mu_{30} = 0.5$ (lower). The two vertical lines bracket the range of observed periods. Note the wide range of mass inflow rates.

Source	P (s)	$\dot{P}$ (s s ⁻¹)	$\dot{M}_{15}$	P_{eq} (s)	t_o (yrs)	μ_{30}	t_{SNR} (yrs)
		$\dot{\Omega}$ (rad s ⁻²)					
1E1048.1-5937	6.44	2×10^{-11}	1	23	7.4×10^3	1.45	
		0.2	64.4	9.3×10^3	2.14		
1E2259+586	6.98	6×10^{-13}	0.5	8.25	5.7×10^4	0.67	
			2	7.4	1.7×10^4	0.54	$< 2 \times 10^4$
			100	6.99	367	3.57	
4U0142+61	8.69	2.1×10^{-12}	1	10.5	2.1×10^4	0.57	
RXJ170849.0	11	2×10^{-11}	12	12.1	1.5×10^3	2.36	
1E1841-045	11.8	4×10^{-11}	3.5	17.5	3.2×10^3	1.96	$< 3 \times 10^3$
AXJ1845-0258	6.97	(7.8×10^{-12}) (-10^{-12})	1	16.8	1.2×10^4	(1)	$< 8 \times 10^3$
SGR1806-20	7.47	8.3×10^{-11}	100	13.8	148	7.9	
SGR1900+14	5.16	1.1×10^{-10}	20	9.1	1.3×10^3	2.2	
RXJ0720.4-3125	8.39	(2.9×10^{-11}) (-2.6×10^{-12})	1	10.8	2×10^4	0.59	
RXJ0420.0-5022	22.7	(8.2×10^{-10}) (-10^{-11})	6.4	7.58	5.27×10^3	3.6	
RQNS in Cas A		(4.7×10^{-11}) (-3×10^{-10})	735	0.99	689	(1)	

Table 1: Model parameters for all six AXPs, for the two SGRs with observed P and $\dot{P}$, the two DTNs with observed periods, and the source in Cas A as a representative RQNS. For the AXPs and SGRs, mass inflow rates chosen are greater than the mass accretion rate inferred from the observed luminosity. For AXJ1845-0258, for which $\dot{P}$ has not been measured, we assume $\mu_{30} = 1$. For the two DTNs and for the RQNS in Cas A, we infer $|\dot{\Omega}|$ from the luminosity. For the DTNs, we assume $P = P_{eq}$, and for the Cas A source $\mu_{30} = 1$ is assumed. Parentheses mark input quantities that are not taken from observations.